

BIOADVANCED
SCIENCE-BASED SOLUTIONS

**PEOPLE
& PETS**

MAY RE-ENTER ONCE
GRASS HAS DRIED




COMPLETE BRAND INSECT KILLER

FOR LAWNS READY-TO-SPREAD GRANULES

SEASON-LONG CONTROL

Starts to Kill in Minutes

***Kills Insects Above
& Below the Ground***



ACTIVE INGREDIENT:	
Lambda-cyhalothrin	0.1%
Other Ingredients	99.9%
Total	100.0%

FOR OUTDOOR USE AROUND THE HOME ONLY
KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional
precautionary statements



Mata todos los tipos de plagas invasores de la lista
Control durante toda la temporada
Mata por contacto

TREATS UP TO
12,500 SQ. FT.
NET WEIGHT 10 lb. (4.5 kg.)



COMPLETE^{BRAND} INSECT KILLER

FOR LAWNS READY-TO-SPREAD GRANULES

*Kills all types of listed lawn damaging insects.
Kills on contact.
Starts to work immediately, lasts all season long.
Prevents lawn damage for a stronger lawn.*

DIRECTIONS FOR USE

READ ENTIRE LABEL BEFORE USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **Keep children and pets off treated area until product has been watered in and lawn is dry. In New York State, this product may not be applied to lawns within 100 feet of a coastal marsh, or stream that drains directly into a coastal marsh.**

GENERAL INFORMATION

For use outdoors, on lawns and as a barrier band treatment around house foundation. Kills ants (including Fire Ants), armyworms, billbugs, chinch bugs, crickets, cutworms, earwigs, European crane flies, fleas, grubs, roaches (German and American), sod webworms, ticks and other insect pests listed in the tables below.

Know Grubs. Stop Grubs.

White grubs are the immature larvae of scarab beetles. These extremely destructive turf pests feed on grass roots just below the soil surface, severing roots and causing lawn grasses to brown and die. Effective white grub control involves reaching these pests at their most vulnerable. Fall treatment targets newly hatched grubs while they're still small and close to the surface.

With heavy infestations, you can roll turf back — just like a rug — to reveal severed roots and feeding grubs. To check your lawn, use a spade to lift a 12-inch-square piece of sod about 3 inches thick. Inspect the soil, thatch and roots closely.

OVERALL LAWN APPLICATION

PESTS	USE RATE	USE INSTRUCTIONS
Ants†, Armyworms*, European Crane Flies, Crickets ^Ø , Cutworms*, Earwigs, Fleas, Lawn moths (sod webworms), Grasshoppers**, Japanese beetle (adults), Pillbugs, Spittle bugs, Mites, Green June Beetles, Leafhoppers [§] , Millipedes, Springtails and Ticks (including Ticks which may transmit Lyme Disease).	Apply at the rate of 0.8 lb per 1,000 sq. ft. (an area 20' x 50') For season long control apply at the rate of 2 lb per 1,000 sq. ft. (an area 20' x 50') Make no more than 6 applications per year at the low use rate or 3 times per year at the highest use rate.	• Apply with a fertilizer spreader (see spreader setting chart). Apply when grass is dry and water the lawn lightly immediately following application. • For season long control of insects above the soil, including Red Imported Fire Ants, apply at higher use rate, as an overall lawn application. • Treat when insects first appear or when you notice lawn damage.
Billbugs, White Grubs (including Japanese beetle grubs, Green June Beetle Grubs, Black dung beetle grubs, Brown dung beetle grubs, Night-flying dung beetle grubs, Asiatic garden beetle grubs, Black Turfgrass Ataenius grubs, European Chafer grubs, May Beetle grubs, June Beetle grubs, Northern Masked Chafer grubs, Oriental Beetle grubs, Southern Masked Chafer grubs), Ataenais spretulus (dung beetles) Hyperodes Weevils, Mole Crickets, Chinch bugs	Apply 1.2 lb. per 1,000 sq. ft. of lawn. Make no more than 6 applications per year at the low use rate or 3 times per year at the highest use rate.	• Water grass completely immediately following application. • Treat when insects first appear or when you notice lawn damage. • For hyperodes weevils , apply in April-May. • For billbug control , apply in May or when activity is first observed or when chewed or brown grass indicates damage from these insects. • For grub control , except on dung beetle, treat any time between late July and early October. • For dung beetle grub control , apply in early spring or early summer (March-July) when damage is first noticed. Immediately following application, completely water grass with ¼ to ½ inch of water. If a second generation occurs, treat again in late summer or early fall.

† Includes Argentine, Florida Carpenter, Black Carpenter, Southern, Field, Allegheny Mound, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance
* Armyworms including: Fall, Yellow-striped and Lawn
Ø Crickets including: House, Common short-tailed, Indies short-tailed, Arizona, Tropical, Variable field, Fall field, Southeastern field, Texas field, Spring field, Northern wood, Western striped, Eastern striped, Sand field, Vocal field, and Southern wood
* Cutworms including: Army, Black, Bronze, Clover, Granulate, Greasy, Green, Spotted, and Variegated
** Grasshoppers including: Banded range, Carolina, Differential, Horse lubber, Panther spotted, Pygmy, Red legged, Southeastern Lubber, Two striped, Alutacea Bird, American Bird, Angular-winged Katydid, Aztec Pygmy
§ Leafhoppers including: Aster, Blunt-nosed, Elm, Mountain, Red-banded, Rice, Rose, Sharp-nosed, Six-spotted, Southern garden, Variegated, Variegated larvae, Virginia creeper, White-banded elm

SPREADER SETTINGS FOR LAWN APPLICATION

NOTE: Spreader settings are approximate. Calibrate your spreader before application since manufacturer's settings can vary among the same model of spreader.

BROADCAST SPREADER	SETTING		
	.8 lb/1,000 sq ft	1.2 lbs/1,000 sq ft	2.0 lbs/1,000 sq ft
Scotts Basic	2.75	3.25	4.25
Scotts Standard	2.75	3.25	4
Scotts Deluxe Edgeguard	2.75	3.25	3.75
Vigoro 10000	4.75	5.25	6.5
Vigoro 15000	4.75	5.25	6.5

DROP SPREADER	SETTING
	2.0 lbs/1,000 sq ft
Scotts Accugreen 1000	3.25
Scotts Accugreen 2000	3.25
Scotts Accugreen 3000	3.25

STORAGE AND DISPOSAL PESTICIDE STORAGE: Reclose container tightly after each use. Store in a cool, dry area which is inaccessible to children and pets.	CONTAINER HANDLING: IF EMPTY: Do not reuse this container. Place in trash or offer for recycling if available. IF PARTLY FILLED: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. Note: Buyer assumes all responsibilities for safety and use not in strict accordance with directions.
---	--

FIRST AID - IF IN EYES: • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice. Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.	For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents) you may contact SafetyCall® International at 1-(877)-229-3724, 24 hours per day, 7 days per week.
---	---

ALMACENAMIENTO Y ELIMINACIÓN ALMACENAMIENTO DEL PESTICIDA: Vuelva a cerrar muy bien el contenedor después de cada uso. Almacenar en u lugar fresco, seco que no esté al alcance de niños y mascotas.	MANIPULACIÓN DEL CONTENEDOR: Si está vacío: No vuelva a usar este contenedor. Colocar en la basura u ofrecer para su reciclado si fuese posible. Si está parcialmente lleno: Llamar a la agencia de eliminación de desechos sólidos para conocer las indicaciones de eliminación. Nunca colocar el producto sin usar en un drenaje de salida o de entrada. Nota: El comprador asume toda la responsabilidad por seguridad y uso del mismo que no esté en estricto acuerdo con las indicaciones de uso.
--	---

PRIMEROS AUXILIOS SI ENTRA EN CONTACTO CON LOS OJOS: • Mantener los ojos abiertos y enjuagar lenta y cuidadosamente con agua durante 15 - 20 minutos. • Quitar los lentes de contacto, si los tuviera, luego de los primeros 5 minutos, continuar lavando. • Llamar al centro de control de envenenamiento o al médico para asesoramiento sobre el tratamiento. Tenga el contenedor o etiqueta del producto con usted cuando llame al centro de control de envenenamiento o al médico, o cuando vaya para recibir tratamiento. Para acceder a información adicional sobre este producto pesticida, (incluyendo dudas de salud, emergencias médica s o incidentes con el pesticida), puede comunicarse con SafetyCall® International al 1-(877) 229-3724, las 24 horas del día, los 7 días de la semana.
--

Apply in three easy steps:

- (1) Determine the size of the job and measure the area to be treated. Using the spreader setting chart, set your spreader to the proper setting.
- (2) Apply product according to labeled rates. Complete coverage is important.
- (3) Water in granules immediately following application to release the active ingredient into the grass and soil. People and pets may re-enter lawn once grass has dried.

NOTE:

- Make no more than 6 applications per year per location at the low use rate (0.8 lb. per 1000 ft²).
- Make no more than 3 applications per year per location at the high use rate (2 lb. per 1000 ft²).
- Do not allow product to run off from target area.

BAND TREATMENT

Around house foundation to control pests normally found around or near foundations of houses, as well as in lawns, and certain of these pests that may enter houses.

PESTS	USE RATE	USE INSTRUCTIONS
Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance), Fleas, Roaches (German and American), Palmetto bugs, Waterbugs, Millipedes, Mites, Silverfish, Sowbugs (Pillbugs), Crickets, Earwigs	Apply at the rate of 0.8 lb per 1,000 sq. ft. (an area 5' x 200') or 1.25 oz. per 100 sq. ft. (5' x 20').	• Apply a 3 to 5 foot band of granules next to the house foundation as well as lawns. Evenly distribute the granules. Water lightly immediately following application. • Treat when insects first appear or when you notice lawn damage.

SPOT TREATMENT

PESTS	USE RATE	USE INSTRUCTIONS
Ants (Argentine, Florida Carpenter, Black Carpenter, Southern, Field, Allegheny Mound, Cornfield, Honey, Pavement, Nuisance)	Mound Treatment: Apply 1/2 teaspoon over and around each ant hill.	Water lightly immediately following application.
Red Imported Fire Ants	Mound Treatment: Apply 1/4 cup over, and 2 ft. around, each fire ant mound.	Immediately following application, water in completely. Apply a minimum of ¼ cup of the product in 1 gallon of water to each mound area. To prevent the disturbance of ants, water gently. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. NOTE: For best results, apply in cool weather, 65°-80° F, or in early morning or late evening hours. Treat new mounds as they appear. Fast acting. Works within 24 hours.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. To protect the environment, **DO NOT** allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS

DECLARACIONES DE PRECAUCIÓN

RIESGOS PARA SERES HUMANOS O ANIMALES DOMÉSTICOS

PRECAUCIÓN - Causa irritación moderada en los ojos. Evitar el contacto con los ojos o ropa. Lavar las manos cuidadosamente con agua y jabón luego de manipularlo y antes de comer, beber, masticar goma de mascar o usar tabaco. No se deben usar utensilios tales como cucharas o tazas medidoras con fines de alimentación luego de usarlos para medir los pesticidas.

RIESGOS AMBIENTALES

Este producto es muy tóxico para los peces y los organismos acuáticos y tóxico para la vida salvaje. Este producto es altamente tóxico para las abejas expuestas al tratamiento directo o residuos en plantas con flores o malezas. **NO** aplicar este producto o permitir que se escurra por malezas o cultivos por nacer si las abejas buscan alimento en esta área. Para proteger al medio ambiente, **NO** permita que el pesticida ingrese o corra por los drenajes de tormenta, drenaje de aguas pluviales, canaletas o aguas superficiales. Aplicar este producto en clima calmo cuando no haya pronóstico de lluvia en las próximas 24 horas para asegurar que el viento o la lluvia no soplen o laven el pesticida fuera del área de tratamiento. Barrer todo producto que quede en caminos, aceras o calles, de nuevo hacia el área tratada del césped o jardín ayudará a evitar que el producto ingrese en cuerpos de agua o sistemas de drenaje.

BioAdvanced® is a registered trademark of SBM Life Science Corp.
Distributed by: SBM Life Science Corp., PO Box 525, Morrisville, NC 27560





COMPLETE INSECT KILLER

FOR LAWNS READY-TO-SPREAD GRANULES